

B7
concl

receptors (LIRs) which bear homology to killer inhibitory receptors (KIRs). These molecules interact with MHC-class I molecules via Ig-like domains and regulate negatively the activation of APC, recruiting an inhibitory signaling molecule, tyrosine phosphatase SHP-1. These data indicate that Ts-induced suppression of APC is based on an active mechanism by up-regulating the expression of a class of inhibitory receptors which transmit negative inhibitory signals in APC. Ts provides an essential regulatory mechanism through which ~~immune tolerance can be achieved.~~

In the title:

On pages 1 and 189, please change the title to read:

B8
Methods for Generating Tolerogenic Antigen Presenting Cells

In the claims:

Please amend the claims as follows:

B9
38. (Amended) A method of generating a tolerogenic antigen presenting cell (APC) which comprises:

- a) contacting an APC with a CD8+CD28- Ts; and
- b) causing overexpression, in the APC of step (a), of mRNA which encodes an inhibitory monocyte inhibitory receptor (MIR), thereby generating a tolerogenic antigen presenting cell (APC).

39. (Amended) The method of claim 38, wherein the inhibitory (MIR) is selected from the group consisting of ILT4 (MIR-10), ILT2 (MIR7), and ILT3.

✓
Please add new claim 70: